

# Experimental detection of the actinospores of *Myxobolus pseudodispar* (Myxosporea: Myxobolidae) in oligochaete alternate hosts

Cs. Székely\*, K. Molnár, E. Eszterbauer, F. Baska

Veterinary Medical Research Institute, Hungarian Academy of Sciences, PO Box 18, 1581 Budapest, Hungary

**ABSTRACT:** The development of *Myxobolus pseudodispar* Gorbunova, 1936, an intracellular myxosporean muscle parasite of the roach *Rutilus rutilus* L., was studied in experimentally infected oligochaetes. In one experiment, uninfected *Tubifex tubifex* Müller and *Limnodrilus hoffmeisteri* (Claparède) were exposed to mature spores of *M. pseudodispar*. Triactinomyxon spores developed both in *T. tubifex* and *L. hoffmeisteri* specimens. Triactinospores were first released from the oligochaetes 76 d after initial exposure. At that time, pansporocysts containing 8 triactinospores were located in the gut epithelium of experimentally infected oligochaetes, but free actinosporean stages were also found in their gut lumen. Each triactinospore had 3 pyriform polar capsules and an elongated cylindrical sporoplasm with 8 secondary cells. The spore body joined the 3 caudal projections with a relatively long style. One of the 3 caudal projections was shorter than the other two. The total length of the triactinospore was on average 206.5 µm.

**KEY WORDS:** *Myxobolus pseudodispar* · Myxosporea · *Rutilus rutilus* · Development · Triactinomyxon stage · *Tubifex tubifex* · *Limnodrilus hoffmeisteri*

## INTRODUCTION

*Myxobolus pseudodispar*, a species described by Gorbunova (1936), is a myxosporean commonly occurring in the skeletal muscles of the roach *Rutilus rutilus* L. in Hungarian lakes, rivers and ponds (Baska 1986). Some aspects of its location and intrapiscine development were also studied by Baska (1986). Nothing was known, however, about its extrapiscine developmental stage.

The extrapiscine phase of a myxosporean was first studied by Wolf & Markiw (1984), who revealed that the extrapiscine development of *Myxobolus cerebralis* Hofer took place in an oligochaete alternate host (*Tubifex tubifex* Müller). Following that pioneering work, several other papers were published to support its finding. These papers (El-Matbouli & Hoffmann 1989, 1993, Ruidisch et al. 1991, Styer et al. 1991, El-Matbouli et al. 1992, Grossheider & Körting 1992, Benajiba & Marques 1993, Kent et al. 1993, Yokoyama

et al. 1993, 1995, Uspenskaya 1995, Trouillier et al. 1996, Yokoyama 1997) revealed that in each case various *Oligochaeta* spp. or *Polychaeta* sp. (Bartholomew et al. 1997) acted as alternate hosts in the development of various myxosporean species.

Recently the life cycles of 7 species from the genera *Myxobolus*, *Thelohanellus* and *Sphaerospora* have also been experimentally studied in our laboratory (El-Mansy & Molnár 1997a,b, El-Mansy et al. 1998a, Székely et al. 1998, Molnár et al. 1999a,b).

The work presented in this paper is part of continuing experimental life cycle studies conducted on the most common myxosporeans of Hungarian fishes. In the experiments reported here, the development of *Myxobolus pseudodispar* was followed in *Tubifex tubifex* and *Limnodrilus hoffmeisteri* alternate hosts.

## MATERIALS AND METHODS

Myxospores of *Myxobolus pseudodispar* were collected from mature muscle cysts of 2 to 5 yr old roach *Rutilus rutilus* specimens seined in Lake Balaton, Hun-

\*E-mail: szekely@novell.vMRI.hu

gary. Pieces of the muscle were cut, compressed between 2 glass plates, and examined for the presence of myxosporean plasmodia. Plasmodia located intracellularly within muscle cells were separated and opened with a needle under a stereomicroscope. Mature spores obtained from plasmodia were identified and counted. A total of 190 000 spores of *M. pseudodispar* were collected.

Oligochaetes *Tubifex tubifex* Müller and *Limnodrilus hoffmeisteri* (Claparède), identified according to Brinkhurst (1963), were collected from the outlet of a duck farm in Szarvas, Hungary, where no fishes live. This parasite-free oligochaete stock containing members of the 2 species in about the same number was transferred to sterilised mud, and propagated in the laboratory in aerated aquaria. The worms were fed on some drops of granulated fish food.

The experiment started on 4 September 1998, when 100 specimens each of *Tubifex tubifex* and *Limnodrilus hoffmeisteri* were placed into a small plastic cup of 500 ml volume, to which spores of *Myxobolus pseudodispar* were added. The calculated spore count per oligochaete was a total of 1000. The temperature of the room during the experiment varied between 20 and 23°C. All dishes were permanently aerated and regularly supplied with fresh water to prevent evaporation and to refresh the water for the oligochaetes. Starting from 42 d post-exposure (p.e.), water from the dishes was filtered weekly through a fine mesh of 21 µm pore size. The filtrates were taken up in a drop of water and examined for the presence of actinospores. On Day 76 p.e. 150 oligochaetes were placed into 2 ml cell-well plates as described by Yokoyama et al. (1991), and

after 1 d of incubation they were examined for the release of actinospores under a stereomicroscope. Triactinospores released by the oligochaetes were examined under a coverslip. They were recorded on videotapes and digitized pictures were taken with the help of the IMAGO® program as described by Székely (1997). Photographs were taken, drawings made and measurements of 47 actinospores recorded. In the description, all measurements are given in µm. The actinosporean stage of *M. pseudodispar* was described using the terminology of Janiszewska (1957) as modified by Lom et al. (1997). After the first release of actinospores live *T. tubifex* and *L. hoffmeisteri* specimens were also examined for actinosporean stages under a compound microscope. After finding the first released triactinospores 2 infected *T. tubifex* and 1 *L. hoffmeisteri* specimen were fixed in 5% glutaraldehyde for 1.5 h and post-fixed in 2% osmium tetroxide for 2 h, washed several times with cacodylate buffer, dehydrated and embedded in Durcupan ACM resin. Semithin sections (0.5 to 1 µm) were made and stained with 0.1% toluidine blue solution.

## RESULTS

### Light microscopy

Actinospores of *Myxobolus pseudodispar* were released from both *Tubifex tubifex* and *Limnodrilus hoffmeisteri*, but no infection was found in the control tanks of the same oligochaete species. Of 150 specimens examined in cell-well plates, 4 *T. tubifex* (5.3%)




Fig. 1. Cross-section of *Tubifex tubifex* infected by *Myxobolus pseudodispar*. The intestine of the oligochaete is heavily infected with pansporocysts (arrowheads) located in the epithelium. Some released pansporocysts are found also in the intestinal lumen (1). Semithin section.  $\times 250$




Fig. 2. Cross-section of *Tubifex tubifex* infected by *Myxobolus pseudodispar*. Enlarged part of Fig. 1. A spansporocyst containing 8 actinospores is located in the gut epithelium. Inside the pansporocyst sporoplasm, polar capsules (arrowhead) and folded membranes of the caudal processes (m) of triactinospores are seen. Semithin section.  $\times 2000$

and 2 *L. hoffmeisteri* (2.7%) proved to be infected. During the experiment, only triactinomyxon-type actinospores were obtained from all infected worms. The release of triactinospores lasted for about 30 d. In infected live oligochaete specimens placed under a coverslip, pansporocysts in the gut epithelium and free spores in the gut lumen of the worms were easily seen under a compound microscope. No infection was found in the control oligochaete stock examined in the same numbers.

### Histology

Pansporocysts of *Myxobolus pseudodispar* were mostly found in the gut epithelium, but pansporocysts and free spores released into the intestinal lumen through the damaged gut epithelium were also seen in semi-thin sections (Fig. 1). Eight triactinospores could be counted in a pansporocyst. In horizontally sectioned triactinospores the spore body with the sporoplasm and the dark-stained polar capsules could also be recognised (Fig. 2) Inside the sporoplasm large, round secondary cells were distinguishable. The free part of the spore body inside the pansporocyst was filled with the folded styles and caudal processes of the actinospores, which could easily be differentiated from the spore body by their pale colour.

**Description of triactinospores.** Triactinospores (Table 1, Figs. 3 & 4) released from the tubificid body and floating in the water were characterised by 3 pyriform polar capsules, a sporoplasm, a style and 3 caudal

processes (Fig. 4). Polar capsules, pyriform in shape, were 5.1 (4–6)  $\mu\text{m}$  in length and 3  $\mu\text{m}$  in width. The cylindrical sporoplasm, 45 (42–48)  $\mu\text{m}$  long and 15.8 (12–18)  $\mu\text{m}$  wide, contained 8 spherical secondary cells 7.6 (6.4–8)  $\mu\text{m}$  in diameter (Fig. 3 inset). The style was relatively long, 157.3 (145–173)  $\mu\text{m}$  in length. Its width measured 15.8 (12.8–18)  $\mu\text{m}$  at the middle. Caudal processes were unequal in length, one of them being shorter. They were slightly bent, tapering toward the end and terminating in a sharply pointed end. The length of the 2 longer caudal processes was 196 (190–204)  $\mu\text{m}$ , that of the shorter one 127.2 (104–144)  $\mu\text{m}$ . Their width at the middle was 13.6 (12–16)  $\mu\text{m}$ .

Table 1. *Myxobolus pseudodispar*. Measurements of the triactinospores (on the basis of 47 triactinospores) in  $\mu\text{m}$

Measure		Mean $\pm$ SD	Minimum	Maximum
Total length		206.5 $\pm$ 15.6	179.2	244
Polar capsules	Length	5.1 $\pm$ 0.5	4	6
	Width	3.0 $\pm$ 0.0	3	3
Sporoplasm	Length	45.0 $\pm$ 2.2	42	48
	Width	15.8 $\pm$ 1.1	12.8	18
Spore body	Length	50.4 $\pm$ 2.4	47	53
	Width	15.8 $\pm$ 1.1	12.8	18
Style	Length	157.3 $\pm$ 14.8	145	173
	Width	15.8 $\pm$ 1.1	12.8	18
Caudal processes				
Longer	Length	196.6 $\pm$ 5.7	190	204
	Width	13.6 $\pm$ 1.7	12	16
Shorter	Length	127.2 $\pm$ 14.8	104	144




Fig. 3. *Myxobolus pseudodispar*. Triactinospores floating in the water.  $\times 500$ . Inset: Spore body of the triactinospore with sporoplasm containing 8 sporoplasm cells and with polar capsules at the apical end.  $\times 1200$

The length from the apical point of polar capsules to the end of sporoplasm (spore body) measured 50.4 (47–53)  $\mu\text{m}$ . The total length from the polar capsules to the end of style (spore body with style) measured 206.5 (179–244)  $\mu\text{m}$ .

#### Differential diagnosis

Both the myxospores and the actinospores of *Myxobolus pseudodispar* can readily be differentiated from those of other species by their asymmetric forms. The triactinomyxon stages of this species resemble other known triactinomyxospores of myxosporean spe-

cies. However, they are distinguishable by the low number of secondary cells in the sporoplasm and by the shorter size of 1 of the caudal processes. The sporoplasm of this species has only 8 large sporoplasm cells and the third caudal process measures only two-thirds of the 2 others.

The triactinomyxons obtained in this experiment are probably identical to one of the triactinomyxon types (T2) detected by El-Mansy et al. (1998b) during a survey of actinospore infection of oligochaetes in a fish farm near Budapest.

#### DISCUSSION

The species *Myxobolus pseudodispar* is a rather common parasite of the roach in Hungary (Molnár 1979, Baska 1986, Székely & Molnár 1996–1997). The validity of the species was questioned by Lom & Dyková (1992), who regarded it as a synonym of *M. cyprini*. In the typical fish hosts (*Cyprinus carpio* and *Rutilus rutilus*) clear differences exist in the spore morphology (Molnár & Kovács-Gayer 1985, Baska 1986), which supports the opinion represented by Gorbunova (1936) and Donec & Shulman (1984); therefore, we regard *M. pseudodispar* as a valid species. In spite of its common occurrence, little is known about its pathogenicity. No studies have been done on the intraoligochaete development of this species. Data obtained in this study show that the intraoligochaete development of this parasite follows the same pattern as described by Wolf & Markiw (1984) for *M. cerebralis*. In the present experiments the development of *M.*




Fig. 4. Schematic drawing of the triactinospore of *Myxobolus pseudodispar*. Bar = 100  $\mu\text{m}$

*pseudodispar* was successfully completed in both *Tubifex tubifex* and *Limnodrilus hoffmeisteri*, in which triactinomyxon-type spores developed. In their shape and size the spores resembled most of the triactinospores known from experimental studies on *Myxobolus* species (*M. cerebralis*, *M. cotti*, *M. carassii*, *M. drjagini*, *M. portucalensis*, *M. hungaricus*), but due to the low number of secondary cells in the sporoplasm and by the shorter length of one of the caudal processes they distinctly differed from the known triactinospores.

According to El-Matbouli et al. (1992), the intra-oligochaete development of *Myxobolus cerebralis*, *M. cotti* and *M. carassii* took 80 to 120 d, while for *M. drjagini*, *M. hungaricus* and *M. portucalensis* El-Mansy & Molnár (1997a,b) and El-Mansy et al. (1998a) reported development times of 91, 102 and 160 d, respectively. The patent period of *M. portucalensis* was somewhat shorter than that of the above-listed species.

The prevalence of infection (5.3% in *Tubifex tubifex* and 2.7% in *Limnodrilus hoffmeisteri*) proved to be relatively low compared to the high levels observed in previous experiments on *Myxobolus drjagini*, *M. portucalensis*, *M. hungaricus* and *M. dispar* (El-Mansy & Molnár 1997a,b, El-Mansy et al. 1998a, Molnár et al. 1999a) in our laboratory. From a single experiment it is impossible to draw conclusions as to whether the alternate hosts, the myxospores or some other factors are responsible for this low prevalence.

Histological studies prove that *Myxobolus pseudodispar* develops in the gut epithelium of the oligochaete alternate host, from where its pansporocysts and spores are released into the lumen and subsequently excreted with the faeces.

Up to now, the intraoligochaete life cycle of about 20 species has been elucidated. However, a technique enabling consistently successful laboratory infection is known only in the case of *Myxobolus cerebralis*. The species *M. pseudodispar* is not ideal for use in experiments, as its relatively small, intracellular plasmodia can be collected and identified only with great difficulty. Reinfection of the fish host with actinospores is also difficult to achieve, as roaches reared infection-free are not yet available. At the same time, the collected myxospores and the obtained actinospores provide excellent material for expanding the molecular biological studies that have already been done on other species (Andree et al. 1999).

**Acknowledgements.** The study was supported by the Hungarian Scientific Research Fund (OTKA, projects No. T 020044 and T 029200). The authors thank Ms Zsuzsa Kis for her help in collecting and culturing tubificids, and are grateful to the Balaton Fisheries Co. for providing fish.

## LITERATURE CITED

- Andree KB, Székely Cs, Molnár K, Gresoviac SJ, Hedrick RP (1999) Relationships among members of the genus *Myxobolus* (Myxozoa: Bivalvidae) based on small subunit ribosomal RNA sequences. *J Parasitol* 85:68–74
- Bartholomew JL, Whipple MJ, Stevens DG, Fryer JL (1997) The life cycle of *Ceratomyxa shasta*, a myxosporean parasite of salmonids, requires a freshwater polychaete as an alternate host. *J Parasitol* 83:859–868
- Baska F (1986) Histological studies on the development of *Myxobolus pseudodispar* Gorbunova, 1936 in the roach (*Rutilus rutilus*). *Acta Vet Hung* 35:251–257
- Benajiba MH, Marques A (1993) The alternation of actinomyxidian and myxosporidian sporadic forms in the development of *Myxidium giardi* (parasite of *Anguilla anguilla*) through oligochaetes. *Bull Eur Assoc Fish Pathol* 13: 100–103
- Brinkhurst RO (1963) A guide for the identification of British aquatic Oligochaeta. *Freshw Biol Assoc Sci Pub* 22:1–55
- Donec S, Shulman SS (1984) Parasitic protozoa. Phylum Cnidosporidia. In: Bauer ON (ed) Key to determination of parasites of freshwater fishes of the USSR, Vol I. Publ House Nauka, Leningrad, p 88–251 (in Russian)
- El-Mansy A, Molnár K (1997a) Extrapiscine development of *Myxobolus drjagini* Achmerov, 1954 (Myxozoa: Myxobolidae) in oligochaete alternative hosts. *Acta Vet Hung* 45:427–438
- El-Mansy A, Molnár K (1997b) Development of *Myxobolus hungaricus* Jacsó, 1940 (Myxozoa: Myxobolidae) in oligochaete alternate hosts. *Dis Aquat Org* 31:227–232
- El-Mansy A, Molnár K, Székely Cs (1998a) Development of *Myxobolus portucalensis* Saraiva & Molnár, 1990 (Myxozoa: Myxobolidae) in the oligochaete *Tubifex tubifex* (Müller). *Syst Parasitol* 41:95–103
- El-Mansy A, Székely Cs, Molnár K (1998b) Studies on the occurrence of actinosporean stages of fish myxosporians in a fish farm of Hungary, with the description of triactinomyxon, raabeia, aurantiactinomyxon and neoactinomyxon types. *Acta Vet Hung* 46:259–284
- El-Matbouli M, Hoffmann RW (1989) Experimental transmission of two *Myxobolus* spp. developing by sporogony via tubificid worms. *Parasitol Res* 75:461–464
- El-Matbouli M, Hoffmann RW (1993) *Myxobolus carassii* Klokacheva, 1914 also requires an aquatic oligochaete, *Tubifex tubifex* as intermediate host in its life cycle. *Bull Eur Assoc Fish Pathol* 13:189–192
- El-Matbouli M, Fischer-Scherl T, Hoffmann RW (1992) Transmission of *Hofereilus carassii* Achmerov, 1960 to goldfish *Carassius auratus* via an aquatic oligochaete. *Bull Eur Assoc Fish Pathol* 12:54–56
- Gorbunova M (1936) Changes in the parasitic fauna of pike and roach according to their age. *Uchenye Zapiski Leningradskogo Ordena Lenina Gosudarstvennogo Universiteta*, No. 7. (Biol Ser) Fasc 3, Problems of Economical Parasitology, Leningrad, p 5–30 (in Russian)
- Grossheider G, Körting W (1992) First evidence that *Hofereilus cyprini* (Doflein, 1898) is transmitted by *Nais* sp. *Bull Eur Assoc Fish Pathol* 12:17–20
- Janiszewska J (1957) Actinomyxidia II. New systematics, sexual cycles, description of new genera and species. *Zool Polon* 8:3–34
- Kent ML, Whitaker DJ, Margolis L (1993) Transmission of *Myxobolus arcticus* Pugachev and Khokhlov, 1979, a myxosporean parasite of Pacific salmon, via a triactinomyxon from the aquatic oligochaete *Stylodrilus heringianus* (Lumbriculidae). *Can J Zool* 71:1207–1211

- Lom J, Dykova I (1992) Protozoan parasites of fishes. In: Series of developments in aquaculture and fisheries science, Vol 26. Elsevier, Amsterdam, p 159–237
- Lom J, McGeorge J, Feist SW, Morris D, Adams A (1997) Guidelines for the uniform characterisation of the actinosporean stages of parasites of the phylum Myxozoa. *Dis Aquat Org* 30:1–9
- Molnár K (1979) Protozoan parasites of fish species indigenous in Hungary. *Parasit Hung* 12:5–8
- Molnár K, Kovács-Gayer É (1985) The pathogenicity and development within the host fish of *Myxobolus cyprini* Doflein, 1898. *Parasitology* 90:549–555
- Molnár K, El-Mansy A, Székely Cs, Baska F (1999a) Development of *Myxobolus dispar* Thelohan, 1895 (Myxosporea: Myxobolidae) in an oligochaete alternate host, *Tubifex tubifex* (Müller). *Folia Parasitol* 46:15–21
- Molnár K, El-Mansy A, Székely Cs, Baska F (1999b) Development of *Sphaerospora renicola* Dykova and Lom, 1982 (Myxosporea: Sphaerosporidae) in oligochaete alternate hosts. *J Fish Dis* 22:1–11
- Ruidisch S, El-Matbouli M, Hoffmann RW (1991) The role of tubificid worms as an intermediate host in the life cycle of *Myxobolus pavlovskii* (Akhmerov, 1954). *Parasitol Res* 7:663–667
- Styer EL, Harrison L, Burtle GJ (1991) Experimental production of proliferative gill disease in channel catfish exposed to a myxozoan-infected oligochaete, *Dero digitata*. *J Aquat Anim Health* 3:288–291
- Székely Cs (1997) Possible applications of video technology and digital image processing in fish parasitology: morphological examination of the groups Apicomplexa and Myxosporea-Actinosporea by video technology. *Bull Eur Assoc Fish Pathol* 17:81–82
- Székely Cs, Molnár K (1996–1997) Preliminary survey of the parasite fauna of some important fish species in the Upper Reservoir of the Kis-Balaton System. *Parasit Hung* 29-30:45–54
- Székely Cs, El-Mansy A, Molnár K, Baska F (1998) Development of *Thelohanellus hovorkai* and *Thelohanellus nikolskii* (Myxosporea: Myxozoa) in oligochaete alternate hosts. *Fish Pathol* 33:107–114
- Trouillier A, El-Matbouli M, Hoffmann R (1996) A new look at the life-cycle of *Hoferellus carassii* in the goldfish (*Carassius auratus auratus*) and its relation to 'kidney enlargement disease' (KED). *Folia Parasitol* 43:173–187
- Uspenskaya AV (1995) Alternation of actinosporean and myxosporean phases in the life cycle of *Zschokella nova* (Myxozoa). *J Euk Microbiol* 42:665–668
- Wolf K, Markiw ME (1984) Biology contravenes taxonomy in the myxozoa: new discoveries show alternation of invertebrate and vertebrate hosts. *Science* 225:1449–1452
- Yokoyama H (1997) Transmission of *Thelohanellus hovorkai* Achmerov, 1960 (Myxosporea: Myxozoa) to common carp *Cyprinus carpio* through the alternate oligochaete host. *Syst Parasitol* 36:79–84
- Yokoyama H, Ogawa K, Wakabayashi H (1991) A new collection method of actinosporeans—a probable infective stage of myxosporeans to fishes—from tubificids and experimental infection of goldfish with the actinosporean, *Raabeia* sp. *Fish Pathol* 26:133–138
- Yokoyama H, Ogawa K, Wakabayashi H (1993) Involvement of *Branchiura sowerbyi* (Oligochaeta: Annelida) in the transmission of *Hoferellus carassii* (Myxosporea: Myxozoa), the causative agent of kidney enlargement disease (KED) of goldfish *Carassius auratus*. *Fish Pathol* 28:135–139
- Yokoyama H, Ogawa K, Wakabayashi H (1995) *Myxobolus cultus* n. sp. (Myxosporea: Myxobolidae) in the goldfish *Carassius auratus* transformed from the actinosporean stage in the oligochaete *Branchiura sowerbyi*. *J Parasitol* 81:446–451

Editorial responsibility: Wolfgang Körting,  
Hannover, Germany

Submitted: May 17, 1999; Accepted: September 3, 1999  
Proofs received from author(s): November 16, 1999