

1 Role of the copepod parasite *Tracheliastes*
2 *maculatus* Kollar, 1836 (Lernaeopodidae) in
3 the common bream (*Abramis brama*) mortality
4 occurring in Lake Balaton, Hungary
5
6
7
8

9 Cs. Székely^{1*}, M. Láng² and K. Molnár¹

11 ¹Veterinary Medical Research Institute, Hungarian Academy of Sciences, H-1143 Budapest, Hungária
12 krt. 21, Hungary; ²Central Agricultural Office, Veterinary Diagnostic Directorate, H-1149 Budapest,
13 Tábornok u. 2, Hungary
14

15
16 **Abstract**

17 *Tracheliastes maculatus* is a copepod crustacean parasite belonging to the family Lernaeopodidae.
18 In the late spring months, around the time of spawning, large numbers of the imago stages of
19 that parasite can be found attached to the scales of common bream in Lake Balaton every year.
20 Due to their bright red colour, the small ulcers formed at the site of parasite attachment are highly
21 conspicuous even to non-specialists, calling attention to the presence of infestation. In 2008, *T.*
22 *maculatus* infestation affecting the bream stock of Lake Balaton proved to be more intensive than
23 usual. The intensity of infestation of the 3- to 5-year-old bream specimens markedly exceeded the
24 usual level of 7–12 parasites per bream, and infestation could be detected also on the 2-year-old
25 bream specimens 12 to 14 cm in length. On the 2-year-old bream specimens, the inflammatory
26 zone of bacterial origin, which was mostly limited to a single scale on breams of bigger body size,
27 involved a larger area, resulting in scaleless areas and extensive superficial erosions developing in
28 the place of the parasitic focus that had become detached together with the attacked scale. When
29 samples from the haemorrhagic areas, from the serum accumulating in the scale pouches and
30 from the kidney were inoculated onto blood agar, colonies of the facultative pathogenic bacterium
31 *Aeromonas hydrophila* grew out. In addition to the sporadic mortality regularly occurring among
32 older fish specimens in the late spring period simultaneously with parasite infestation, this year
33 mortality could be observed also in the younger age groups of bream.

33 **Introduction**

34 The copepod crustacean *Tracheliastes maculatus*
35 Kollar, 1836 belonging to the Lernaeopodidae
36 family and parasitic on the scales of common
37 bream (*Abramis brama*) calls attention to itself
38 with the interesting clinical signs it causes,
39 rather than with its common occurrence.
40 These greenish, transparent arthropods about

1–1.5 cm in length attach to the scales of fish
with the bulla of their specially elongated
second maxilla, thus securing their permanent
adherence to the host. All of the parasite
specimens attached in this way and visible
also with the unaided eye are females, and
the male specimen of very small size, pressed

* Corresponding author's email: szekely@vmri.hu

close against the female's body, can be detected only on closer inspection. During infestation with *T. maculatus* specimens, which occurs primarily in common bream and less often in white bream and roach, a haemorrhage due to bacterial superinfection develops around the site of parasite attachment, under the scales, in an area involving either a single scale or 3–5 scales, resulting in the development of highly conspicuous red spots on the body of the fish. Females of these parasitic copepods colonising the body surface of common bream by adhering to the scales can be observed in Lake Balaton every year. These crustaceans cause a parasitosis resembling *Lernaea cyprinacea* infestation well known from fish farms. This disease regularly occurs in Lake Balaton in late May or early June, simultaneously with the spawning of common bream, but the severity of the losses caused by it vary from year to year. The appearance of *T. maculatus* infestation in Lake Balaton regularly coincides with a low-level mortality among common bream. The adjective 'maculatus' in the name of the parasitic copepod *T. maculatus* obviously refers to the haemorrhages caused by the bacterial complications of parasitic infestation. The occurrence of the parasite in Lake Balaton and the clinical signs produced by it were first described by Geyer (1939). According to Gussev *et al.* (1987) this parasite is common in the former Soviet Union, but it occurs in Great Britain and in the Netherlands as well (Boxhall and Frear, 1990; Walker *et al.*, 2006). This copepod, which is primarily specialized for the common bream, is a frequent cause of mortality in lakes where the common bream has substantial dominance within the fish population. Such cases were reported by Grabda and Grabda (1958) as

well as Piasecki (1989; cited by Piasecki and Avenant-Oldewage, 2008). The damage done by *T. maculatus* in Lake Balaton was reported by Molnár (1965). Another species, *T. polycolpus* Nordmann, 1832 commonly occurs on fluvial fish species in Hungary; however, this latter species adheres to the base of the fins rather than to the scales (Molnár, 1966).

Material and methods

During a multiple-year survey aimed at the regular monitoring of parasitic infections of Lake Balaton fishes, a few dead common bream specimens were noted in the northern region of the lake, around Tihany, in mid-May 2008. One week later, on the southern shore of the lake, in Balatonszemes higher numbers of dead common bream specimens were recorded. Some moribund 2-year-old common bream, on the body surface of which the red spots and the scaleless areas could already be seen while they were still in the water, were collected with a hand net from the water close to the surface. A higher number of bream specimens showing the clinical signs of the disease were caught with a dragnet. The fish were placed into plastic bags and transported to the laboratory, where the parasites and the lesions caused by them were examined by microscopy and the secondary bacterial infections were studied by inoculation of samples onto blood agar media. Tissue samples from the damaged parts of the skin were fixed in Bouin's solution, embedded in paraffin wax, cut to 4–5 µm sections, and stained with haematoxylin and eosin.

Results and Discussion

During regular fish health monitoring surveys conducted in Lake Balaton in 2008,

1
2
3
4
5
6
7
8
9
10
11
12
13
14
15
16
17
18
19
20
21
22
23
24
25
26
27
28
29
30
31
32
33
34
35
36
37
38
39
40

1 in May and June as in the previous years. *T.*
2 *maculatus* infestations were observed on the
3 scales of common bream accompanied by the
4 appearance of red spots, and simultaneously a
5 low-level common bream mortality was noted,
6 which is known to occur regularly at that time.
7 A unique feature of this infestation was that,
8 unlike the previous years, in 2008 an intensive
9 *T. maculatus* infestation was detected not only
10 in the several years old bream specimens but
11 also in the younger age groups of common
12 bream, and at the same time the mortality rate
13 also exceeded the usual level recorded in the
14 previous years.

15
16 A further interesting feature of the infestation
17 observed in 2008 was that, in addition to
18 the usual inflammation mostly restricted
19 to a single scale or extending only to the
20 immediate surroundings of the attachment
21 site, in the younger bream specimens an
22 erosion extending to large areas of the body
23 surface developed. In the 3- to 5-year-old age
24 group the usual clinical signs, characterised
25 by a ring of inflammation surrounding the
26 attached copepods, were diagnosed. In these
27 areas, at the centre of the haemorrhagic ring
28 *T. maculatus* females, less conspicuous because
29 of their greenish colour, protruded from the
30 affected scale. In these fish groups, as many
31 as 5 to 30 spots indicative of infestation were
32 found on the bream specimens that were
33 infested by the copepods but did not show
34 signs of disease. In the infested 2-year-old
35 common bream specimens 12–14 cm in length,
36 a few copepods attached to the scales were still
37 visible in the areas showing less pronounced
38 inflammation, while on the majority of the body
39 surface only the pathological consequences of
40 infestation could be observed. As a result of

the damage done by the parasite, scaleless
areas of varying size and the appearance of
haemorrhagic-inflammatory foci were seen.
In addition to the direct damage done to the
scales, the bacterial infection developing in
the disintegrated tissues was also a major
factor in the pathogenesis of the disease.
The bright-coloured ulcers were well visible
even with the unaided eye on the body of fish
swimming close to the water surface, often
floating apathetically on their sides or with
their belly up (Figure 1). On the moribund
or dead fish drifted to the shore, there were
confluent haemorrhages and scaleless areas of
substantial size on the sides of the body. The
remaining scales surrounding the erosions had
a ruffled appearance and a serosanguineous
exudate had accumulated under them.

When samples from the haemorrhagic areas,
from the serum accumulating in the scale
pouches and from the kidney were inoculated
onto blood agar, colonies of the facultative
pathogenic bacterium *Aeromonas hydrophila*
grew out. In histological sections accumulation
of serum in the scale pouches was found
(Figure 2), the multilayered connective tissue
of the dermis became disintegrated, and
necrotic cells and tissue debris could be seen
in the serum. As a sign of regeneration, in the
scaleless areas a relatively intact epithelium
developed, which obscured the injured
epithelial cells and the haemorrhagic areas
(Figure 3). In this granulation tissue, red blood
cells, necrotic epithelial cells and occasionally
flint algae could also be observed.

When receiving news of the low-level common
bream mortalities regularly occurring in May,
members of the Fish Pathology Research



Figure 1. Unlike the previous years, in 2008 we detected intensive *T. maculatus* infestation not only in the several years old common bream age groups but also on the younger, 2- to 3-year-old bream specimens (arrows).




Figure 2. As a result of infection, serum (s) accumulates in the scale pouches and the scales assume a "ruffled" appearance.

1
2
3
4
5
6
7
8
9
10
11
12
13
14
15
16
17
18
19
20
21
22
23
24
25
26
27
28
29
30
31
32
33
34
35
36
37
38
39
40




Figure 3. As a sign of regeneration, in the scaleless areas a relatively intact epithelium (e) containing also sensory cells (arrows) appears, which obscures the injured epithelial cells (ie) and the haemorrhagic areas (h).

Team of the Hungarian Academy of Sciences acknowledge this as an unalterable event indicating that the parasitic copepod *T. maculatus* has again exerted its harmful activity. The time of occurrence is also natural, since – as established by Piasecki (1989) – the Cyclops-like larval stages of *T. maculatus* are free-living constituents of the plankton, and its imagos colonise the scales of fish only in the late spring period. Having adhered to the scales, the females undergo substantial transformation. Staying on the scales for about one month, they produce eggs and then die. It is interesting that Gussev *et al.* (1987) mentions the occurrence of *T. maculatus* infestation in the white bream and less often also in other cyprinids, while during this survey the parasite was not detected either on the white bream or on other cyprinid fishes.

As regards the mortality caused by *T. maculatus* infestation, it was unusual that the mortality was observed among young common bream, as until then the mortality caused by this parasite in younger age groups of common bream had not been a typical finding prior to 2008. The sporadic mortalities reported earlier had mostly been restricted to the older age groups of common bream. On the older common bream specimens caught in 2008 the presence of the usual number of *T. maculatus* copepods was noted; however, in addition to the local punctiform haemorrhages involving single scales, the occurrence of confluent haemorrhages extending to larger scaly areas was more frequently seen also on these fish. The number of deaths in these age groups of common bream roughly corresponded to that observed in the previous years. However, the non-negligible mortality rate observed

among the younger, 2- to 3-year-old common bream specimens was a novel feature. By laboratory examination, the accumulation of a serosanguineous exudate under the scales was detectable also in the affected areas still covered with scales. On some scales traces of attachment of copepods that had adhered to the scales but subsequently fallen off was noted; however, the copepods presumably fell off the body surface of these smaller-sized fish together with the scale serving as their site of attachment. All the affected fish transported to the laboratory died in the aquarium within a few days. Both the clinically affected fish and those free from clinical signs were pathologically emaciated. Although the bream is a less fleshy fish even under normal conditions, the body condition of the specimens caught during this survey was poorer than that typical of the Lake Balaton bream. It was also striking that younger bream specimens could be caught using a dragnet near the shoreline as previously it was only possible to catch young bream specimens only from the open waters or from the deep waters of harbours in the autumn or early spring. During a survey conducted in 2009, mortality of the younger bream age groups was not observed, and even among the older age groups only the periodic bream mortality typically found in the earlier years was recorded.

It follows from the biology of this crustacean parasite that its attachment to the body surface of fish requires strong scales. As opposed to *L. cyprinacea*, which fixes its anchors deep in the muscles beneath the pierced scale, the attachment of *T. maculatus* is limited to a single scale. It is presumed that in the year 2008, because of their presence in increased

numbers, *T. maculatus* parasites attacked also the smaller bream specimens but could not permanently establish themselves on them and, in the majority of cases, became detached from the fish together with the attacked scale. The scale injury caused by the parasite increased the chances of bacterial invasion. The limited common bream mortality, typically occurring in Lake Balaton in the period of warming up every year, is partly attributable to the damage done by *T. maculatus* and to secondary *A. hydrophila* infection through the wounds created by the female copepods attached to the scales. The direct cause of death is clearly the septicaemia caused by the facultative pathogenic bacterium *A. hydrophila*, but *T. maculatus* infestation opening the gate for this bacterial infection definitely plays a primary role in the pathogenesis of the disease.

In addition to the pathogen and the environment, the development of the disease greatly depends on the natural resistance of the fish as well. While the continuous improvement of the water quality of Lake Balaton is very pleasing, it is expected to be accompanied by a decrease in the abundance of food organisms available for fish. It was known already from studies conducted in previous years that the growth rate of common bream in Lake Balaton was substantially inferior to that of common bream specimens living in the Kis-Balaton water reservoir, in Lake Velence and in the river Danube more abundant in natural food (Pénzes, 1966; Specziár *et al.*, 1997; Specziár and Tölg, 2000; Specziár and Rezsü, 2009). The feeding opportunities of the common bream stock in Lake Balaton cannot be considered optimal even at present. The increased intensity of

1
2
3
4
5
6
7
8
9
10
11
12
13
14
15
16
17
18
19
20
21
22
23
24
25
26
27
28
29
30
31
32
33
34
35
36
37
38
39
40

1 *T. maculatus* infestation in 2008 may be due
 2 to the conditions favouring the growth of
 3 the parasite and to the different reaction of
 4 younger common bream age groups to the
 5 infestation.

7 **Acknowledgements**

8 This work was supported by the Hungarian
 9 Scientific Research Fund (OTKA), project
 10 No. K 71837. We thank Györgyi Ostoros
 11 for preparing the histological sections. The
 12 help of our colleagues Gábor Cech and
 13 Olga Kelemen (Veterinary Medical Research
 14 Institute, Hungarian Academy of Sciences),
 15 as well as András Specziár and Géza Dobos
 16 (Balaton Limnological Research Institute of
 17 the Hungarian Academy of Sciences) in the
 18 fish collections is gratefully acknowledged.

20 **References**

21 Boxhall GA and Frear PA (1990). *Tracheliastes*
 22 *maculatus* (Kollar, 1936) (Crustacea,
 23 Lernaepodidae), a new species for Britain.
 24 *Journal of Fish Biology* **37**, 489–491.

25 Geyer F (1939). Über parasitische
 26 Lernaepodiden von Balaton Fischen
 27 (Ungarn). *Zoologische Anzeiger* **127**, 145–159.

28 Grabda E and Grabda J (1958). Tracheliastosis
 29 in the common bream, *Abramis brama* L. in
 30 Lake Jamno. *Zoologica Poloniae* **8**, 325–334.

31 Gussev AV, Poddubnaya AV and Avdeyeva
 32 VV (1987). Arthropoda Knidosporidii
 33 (Cnidosporidia). In: **“Key to the Parasites
 34 of Freshwater Fishes of the USSR”** (O. N.
 35 Bauer, Ed.). pp. 378–532. Nauka, Leningrad
 36 (in Russian).

37 Molnár K (1965). On a parasitic copepod
 38 from Lake Balaton. *Halászat* **11**, 151 (in
 39 Hungarian).

40 Molnár K (1966). Cohabitation of a parasitic

copepod and ciliates. *Halászat* **12**, 111 (in
 Hungarian).

Pénzes B (1966). Adatok a balatoni dévérkeszeg
 (*Abramis brama* L.) növekedéséhez [Data on the
 growth of bream in Lake Balaton]. In: **“Annales
 Instituti Biologici (Tihany) Hungaricae
 Academiae Scientiarum”** [Annals of the
 Balaton Limnological Research Institute of the
 Hungarian Academy of Sciences (Tihany)],
 1966, Volume **33**, pp. 173–176. ISSN 1215-
 0584.

Piasecki W (1989). Life cycle of *Tracheliastes*
maculatus Kollar, 1835 (Copepoda,
 Siphonostomatoida, Lernaepodidae).
Wiadomosci Parasitologiczne **35**, 187–245.

Piasecki W and Avenant-Oldewage A (2008).
 Diseases caused by Crustacea. In: **“Fish
 diseases 2”** (Eiras, J. C. et al., Eds). Science
 Publishers Enfield (NH) **18**, 1115–1200.

Specziár A and Rezsű ET (2009). Feeding
 guides and food resource partitioning in a lake
 fish assemblage: an ontogenetic approach.
Journal of Fish Biology **75**, 247–267.

Specziár A and Tölg L (2000). Investigations
 of the bream stock of Lake Balaton. *Halászat*
93 (3), 135–144.

Specziár A, Tölg L and Bíró P (1997). Feeding
 strategy and growth of cyprinids in the littoral
 zone of Lake Balaton. *Journal of Fish Biology* **51**,
 1109–1124.

Walker PD, Abbink W, Van der Velde G
 and Bonga SEW (2006). A new record of
Tracheliastes maculatus Kollar, 1835 (Copepoda,
 Siphonostomatoida, Lernaepodidae) on
 common bream (*Abramis brama* (L., 1758)) in
 the Netherlands. *Crustaceana* **79**, 1015–1019.